

## REFERENCIAS

1. Three-Dimensional Terminology of Spinal Deformity | Scoliosis Research Society [Internet]. [citado 9 de marzo de 2019]. Disponible en: <https://www.srs.org/professionals/online-education-and-resources/glossary/three-dimensional-terminology-of-spinal-deformity>
2. Kalichman L, Kendelker L, Bezalel T. Bracing and exercise-based treatment for idiopathic scoliosis. *J Bodyw Mov Ther.* 2016; 20 (1): 56-64.
3. Hresko MT, Talwalkar V, Schwend R. Early Detection of Idiopathic Scoliosis in Adolescents: *J Bone Jt Surg.* 2016; 98 (16).
4. Peters JR, Chandrasekaran C, Robinson LF, Servaes SE, Campbell RM, Balasubramanian S. Age- and gender-related changes in pediatric thoracic vertebral morphology. *Spine J.* 2015; 15 (5): 1000-20.
5. Negrini S, Donzelli S, Aulisa AG, Czaprowski D, Schreiber S, de Mauroy JC, et al. 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis Spinal Disord.* 2018; 13 (1).
6. Balioglu MB, Aydin C, Kargin D, Albayrak A, Atici Y, Tas SK, et al. Vitamin-D measurement in patients with adolescent idiopathic scoliosis. *J Pediatr Orthop Part B.* 2017; 26 (1): 48-52.
7. Goździalska A, Jaśkiewicz J, Knapik-Czajka M, Drąg J, Gawlik M, Cieśla M, et al. Association of Calcium and Phosphate Balance, Vitamin D, PTH, and Calcitonin in Patients With Adolescent Idiopathic Scoliosis. *Spine.* 2016; 41 (8): 693-7.

8. Mayes T, Anadio JM, Sturm PF. Prevalence of Vitamin D Deficiency in Pediatric Patients With Scoliosis Preparing for Spinal Surgery. *Spine Deform.* 2017; 5 (6): 369-73.
9. Dantas DDS, De Assis SJC, Baroni MP, Lopes JM, Cacho EWA, Cacho RDO, et al. Klapp method effect on idiopathic scoliosis in adolescents: blind randomized controlled clinical trial. *J Phys Ther Sci.* 2017; 29 (1): 1-7.
10. Iunes DH, Cecílio MBB, Dozza MA, Almeida PR. Quantitative photogrammetric analysis of the klapp method for treating idiopathic scoliosis. *Braz J Phys Ther.* 2010; 14 (2): 133-40.
11. Schreiber S, Parent EC, Khodayari Moez E, Hedden DM, Hill DL, Moreau M, et al. Schroth Physiotherapeutic Scoliosis-Specific Exercises Added to the Standard of Care Lead to Better Cobb Angle Outcomes in Adolescents with Idiopathic Scoliosis - an Assessor and Statistician Blinded Randomized Controlled Trial. *PloS One.* 2016 ;11 (12): e0168746.
12. Kwan KYH, Cheng ACS, Koh HY, Chiu AYY, Cheung KMC. Effectiveness of Schroth exercises during bracing in adolescent idiopathic scoliosis: results from a preliminary study-SOSORT Award 2017 Winner. *Scoliosis Spinal Disord.* 2017; 12: 32.
13. Park J-H, Jeon H-S, Park H-W. Effects of the Schroth exercise on idiopathic scoliosis: a meta-analysis. *Eur J Phys Rehabil Med.* 2018; 54 (3): 440-9.
14. Romano M, Negrini A, Parzini S, Tavernaro M, Zaina F, Donzelli S, et al. SEAS (Scientific Exercises Approach to Scoliosis): a modern and effective evidence based approach to physiotherapeutic specific scoliosis exercises. *Scoliosis.* 2015; 10: 3.

15. Negrini A, Negrini MG, Donzelli S, Romano M, Zaina F, Negrini S. Scoliosis-Specific exercises can reduce the progression of severe curves in adult idiopathic scoliosis: a long-term cohort study. *Scoliosis*. 2015; 10: 20.
16. Białek M. Conservative treatment of idiopathic scoliosis according to FITS concept: presentation of the method and preliminary, short term radiological and clinical results based on SOSORT and SRS criteria. *Scoliosis*. 2011; 6: 25.
17. Białek M. Mild Angle Early Onset Idiopathic Scoliosis Children Avoid Progression Under FITS Method (Functional Individual Therapy of Scoliosis): *Medicine (Baltimore)*. 2015; 94 (20): e863.
18. Philippe Souchard. RPG. El método. 1a. Barcelona (España): Elsevier Masson; 2012.
19. Amorim CSM de, Gracitelli MEC, Marques AP, Alves VLDS. Effectiveness of global postural reeducation compared to segmental exercises on function, pain, and quality of life of patients with scapular dyskinesis associated with neck pain: a preliminary clinical trial. *J Manipulative Physiol Ther*. 2014; 37 (6): 441-7.
20. Lozano-Quijada C, Poveda-Pagán EJ, Segura-Heras JV, Hernández-Sánchez S, Prieto-Castelló MJ. Changes in Postural Sway After a Single Global Postural Reeducation Session in University Students: A Randomized Controlled Trial. *J Manipulative Physiol Ther*. 2017; 40 (7): 467-76.
21. Dupuis S, Fortin C, Caouette C, Leclair I, Aubin C-É. Global postural re-education in pediatric idiopathic scoliosis: a biomechanical modeling and analysis of curve reduction during active and assisted self-correction. *BMC Musculoskelet Disord*. 2018; 19.

22. Murrell GA, Coonrad RW, Moorman CT, Fitch RD. An assessment of the reliability of the Scoliometer. *Spine*. 1993; 18 (6): 709-12.
23. Sapkas G, Papagelopoulos PJ, Kateros K, Koundis GL, Boscainos PJ, Koukou UI, et al. Prediction of Cobb angle in idiopathic adolescent scoliosis. *Clin Orthop*. 2003; (411): 32-9.
24. Matamalas A, D'Agata E, Sanchez-Raya J, Bago J. Trunk appearance perception scale for physicians (TAPS-Phy) - a valid and reliable tool to rate trunk deformity in idiopathic scoliosis. *Scoliosis Spinal Disord*. 2016; 11 (1).
25. Thielsch MT, Wetterkamp M, Boertz P, Gosheger G, Schulte TL. Reliability and validity of the Spinal Appearance Questionnaire (SAQ) and the Trunk Appearance Perception Scale (TAPS). *J Orthop Surg*. 2018; 13 (1).
26. Horne JP, Flannery R, Usman S. Adolescent idiopathic scoliosis: diagnosis and management. *Am Fam Physician*. 2014; 89 (3): 193-8.
27. Hammer O, Harper DAT, Ryan PD. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*. 2001; 4 (9): 9.
28. Springate SD. The effect of sample size and bias on the reliability of estimates of error: a comparative study of Dahlberg's formula. *Eur J Orthod*. 2012; 34 (2): 158-63.
29. Klingenberg C. Analyzing Fluctuating Asymmetry with Geometric Morphometrics: Concepts, Methods, and Applications. *Symmetry*. 2015; 7 (2): 843-934.
30. Alves de Araújo ME, Bezerra da Silva E, Bragade Mello D, Cader SA, Shiguemi Inoue Salgado A, Dantas EHM. The effectiveness of the Pilates method: reducing the

- degree of non-structural scoliosis, and improving flexibility and pain in female college students. *J Bodyw Mov Ther.* 2012; 16 (2): 191-8.
31. Kim G, HwangBo P. Effects of Schroth and Pilates exercises on the Cobb angle and weight distribution of patients with scoliosis. *J Phys Ther Sci.* 2016; 28 (3): 1012-5.
  32. Aulisa AG, Giordano M, Falciglia F, Marzetti E, Poscia A, Guzzanti V. Correlation between compliance and brace treatment in juvenile and adolescent idiopathic scoliosis: SOSORT 2014 award winner. *Scoliosis.* 2014; 9 (1): 6.
  33. Konieczny MR, Hieronymus P, Krauspe R. Time in brace: where are the limits and how can we improve compliance and reduce negative psychosocial impact in patients with scoliosis? A retrospective analysis. *Spine J Off J North Am Spine Soc.* 2017; 17 (11): 1658-64.
  34. Rivett L, Rothberg A, Stewart A, Berkowitz R. The relationship between quality of life and compliance to a brace protocol in adolescents with idiopathic scoliosis: a comparative study. *BMC Musculoskelet Disord.* 2009; 10 (1): 5.
  35. Paolucci T, Piccinini G, Iosa M, Piermattei C, De Angelis S, Zangrando F, et al. The importance of trunk perception during brace treatment in moderate juvenile idiopathic scoliosis: What is the impact on self-image? *J Back Musculoskelet Rehabil.* 2017; 30 (2): 203-10.